

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP05/002566

International filing date: 10 February 2005 (10.02.2005)

Document type: Certified copy of priority document

Document details: Country/Office: EP
Number: 04356017.6
Filing date: 12 February 2004 (12.02.2004)

Date of receipt at the International Bureau: 29 March 2005 (29.03.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

10. 02. 2005

**Europäisches
Patentamt****European
Patent Office****Office européen
des brevets****Bescheinigung****Certificate****Attestation**

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

04356017.6

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 04356017.6
Demande no:

Anmeldetag:
Date of filing: 12.02.04
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Bayer CropScience S.A.
16 rue Jean-Marie-Leclair
69009 Lyon
FRANCE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Fungicidal composition comprising a pyridylethylbenzamide derivative and a
compound capable of inhibiting the spores germination or mycelium growth by
acting on different metabolic routes

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A01N1/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

Fungicidal composition comprising a pyridylethylbenzamide derivative and a compound capable of inhibiting the spores germination or mycelium growth by acting on different metabolic routes

5

The present invention relates to novel fungicide compositions comprising a pyridylethylbenzamide derivative and a compound capable of inhibiting the spores germination or mycelium growth by acting on different metabolic routes. The present invention also relates to a method of combating or controlling phytopathogenic fungi by applying at a locus infested or liable to be infested such a composition.

10

International patent application WO 01/11965 generically discloses numerous pyridylethylbenzamide derivatives. The possibility of combining one or more of these numerous pyridylethylbenzamide derivatives with known fungicidal products to develop a fungicidal activity is disclosed in general terms, without any specific example or biological data.

15


It is always of high-interest in agriculture to use novel pesticidal mixtures showing a synergistic effect in order notably to avoid or to control the development of resistant strains to the active ingredients or to the mixtures of known active ingredients used by the farmer while minimising the doses of chemical products spread in the environment and reducing the cost of the treatment.

20

We have now found some novel fungicidal compositions which possess the above mentioned characteristics.

25

Accordingly, the present invention relates to a composition comprising :
a) a pyridylethylbenzamide derivative of general formula (I)



in which :

- p is an integer equal to 1, 2, 3 or 4;
- q is an integer equal to 1, 2, 3, 4 or 5;
- each substituent X is chosen, independently of the others, as being halogen, alkyl or haloalkyl;

30

- each substituent Y is chosen, independently of the others, as being halogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, amino, phenoxy, alkylthio, dialkylamino, acyl, cyano, ester, hydroxy, aminoalkyl, benzyl, haloalkoxy, halosulphonyl, halothioalkyl, alkoxyalkenyl, alkylsulphonamide, nitro, alkylsulphonyl, phenylsulphonyl or benzylsulphonyl;
- as to the N-oxides of 2-pyridine thereof;
- and
- b) a compound capable of inhibiting the spores germination or mycelium growth by acting on different metabolic routes;
- in a (a) / (b) weight ratio of from 0.01 to 20.

In the context of the present invention :

- halogen means chlorine, bromine, iodine or fluorine;
- each of the alkyl or acyl radicals present in the molecule contains from 1 to 10 carbon atoms, preferably from 1 to 7 carbon atoms, more preferably from 1 to 5 carbon atoms, and may be linear or branched;
- each of the alkenyl or alkynyl radicals present in the molecule contains from 2 to 10 carbon atoms, preferably from 2 to 7 carbon atoms, more preferably from 2 to 5 carbon atoms, and may be linear or branched.

The composition according to the present invention provides a synergistic effect. This synergistic effect allows a reduction of the chemical substances spread into the environment and a reduction of the cost of the fungal treatment.

In the context of the present invention, the term "synergistic effect" is defined by Colby according to the article entitled "Calculation of the synergistic and antagonistic responses of herbicide combinations" Weeds, (1967), 15, pages 20-22.

The latter article mentions the formula:

$$E = x + y - \frac{x * y}{100}$$

in which E represents the expected percentage of inhibition of the disease for the combination of the two fungicides at defined doses (for example equal to x and y respectively), x is the percentage of inhibition observed for the disease by the compound (I) at a defined dose (equal to x), y is the percentage of inhibition observed for the disease by the compound (II) at a defined dose (equal to y). When the percentage of inhibition observed for the combination is greater than E, there is a synergistic effect.

The composition according to the present invention comprises a pyridylethylbenzamide derivative of general formula (I). Preferably, the present invention relates to a composition comprising a pyridylethylbenzamide derivative of general formula (I) in which the different characteristics may be chosen alone or in combination as being :

- as regards p, p is 2;
- as regards q, q is 1 or 2. More preferably, q is 2;
- as regards X, X is chosen, independently of the others, as being halogen or haloalkyl. More preferably, X is chosen, independently of the others, as being a chlorine atom or a trifluoromethyl group;
- as regards Y, Y is chosen, independently of the others, as being halogen or haloalkyl. More preferably, Y is chosen, independently of the others, as being a chlorine atom or a trifluoromethyl group;

More preferably, the pyridylethylbenzamide derivative of general formula (I) present in the composition of the present invention is :

- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (compound 1);
- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-iodobenzamide (compound 2); or
- N-{2-[3,5-dichloro-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (compound 3).

Even more preferably, the pyridylethylbenzamide derivative of general formula (I) present in the composition of the present invention is N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (compound 1).

25

The composition according to the present invention comprises a compound capable of inhibiting the spores germination or mycelium growth by acting on different metabolic routes. Preferably, the present invention relates to a composition comprising a compound capable of inhibiting the spores germination or mycelium growth by acting on different metabolic routes selected from dicarboximide derivatives, phthalimide derivatives, 2-butoxy-6-iodo-3-propyl-benzopyran-4-one, 2,6-dichloro-N-{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl}benzamide, (Z)-N-[α -(cyclopropylmethoxyimino)-2,3-difluoro-6-(trifluoromethyl)benzyl]-2-phenylacetamide, benthialvalicarb, chlorothalonil, copper hydroxide, copper oxychloride, copper sulfate, copper sulfate (tribasic), cuprous oxide, cymoxanil, diclomezine, dichlofluanid, dithianon, dimethomorph, dodine, ethaboxam,

35

fenpiclonil, fentin, ferbam, fluazinam, fludioxonil, flusulfamide, guazatine, iminoctadine, mancozeb, mancozeb, maneb, metiram, methasulfocarb, nabam, nickel bis(dimethyldithiocarbamate), iprovalicarb, oxine-copper, propamocarb, propineb, quinoxifen, sulfur, silthiofam, thiram, tolylfluanid, triazoxide, zineb, ziram, phosphorous acid and fosetyl-Al. 2-butoxy-6-iodo-3-propyl-benzopyran-4-one, 2,6-dichloro-N-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl]benzamide, chlorotalonil, iprovalicarb, mancozeb, propamocarb and fosetyl-Al are preferred.

According to the present invention, dicarboximide derivatives may for example be chlozolate, iprodione, procymidone or vinclozolin. Iprodione and are still preferred.

According to the present invention, phthalimide derivatives may for example be captan, captan, folpet, thiochlorfenphim. Captan and folpet are preferred.

The composition according to the present invention comprises (a) at least a pyridylethylbenzamide derivative of general formula (I) and (b) a compound capable of inhibiting the spores germination or mycelium growth by acting on different metabolic routes in an (a) / (b) weight ratio of from 0.01 to 20; preferably of from 0.05 to 10; even more preferably, of from 0.1 to 5.

The composition of the present invention may further comprise at least one other fungicide active ingredient (c).

The fungicidal active ingredient (c) may be selected from azaconazole, azoxystrobin, (Z)-N-[α -(cyclopropylmethoxyimino)-2,3-difluoro-6-(trifluoromethyl)benzyl]-2-phenylacetamide, benalaxyl, benomyl, benthiavalicarb, biphenyl, bitertanol, blastidicidin-S, boscalid, borax, bromuconazole, bupirimate, sec-butylamine, calcium polysulfide, captan, captan, carbendazim, carboxin, carpropamid, chinomethionat, chlorothalonil, chlozolate, copper hydroxide, copper octanoate, copper oxychloride, copper sulfate, cuprous oxide, cymoxanil, cyproconazole, cyprodinil, dazomet, debacarb, dichlofluanid, dichlorophen, diclocymet, diclomezine, dicloran, diethofencarb, difenoconazole, difenzoquat metilsulfate, difenzoquat, diflumetorim, dimethirimol, dimethomorph, diniconazole, dinobuton, dinocap, diphenylamine, dithianon, dodemorph, dodemorph acetate, dodine, edifenphos, epoxiconazole, ethaboxam, ethirimol, ethoxyquin, etridiazole, famoxadone, fenamidone, fenarimol, fenbuconazole, fenfuram, fenhexamid, fenpiclonil, fenoxanil, fenpropidin, fenpropimorph, fentin, fentin hydroxide, fentin acetate, ferbam, ferimzone, fluazinam, fludioxonil, fluoroimide, fluoroimide, fluoroimide, fluoroimide.

fluquinconazole, flusilazole, flusulfamide, flutolanil, flutriafol, folpet, formaldehyde, fosetyl, fosetyl-aluminium, fuberidazole, furalaxyl, furametpyr, guazatine, guazatine acetates, hexachlorobenzene, hexaconazole, 8-hydroxyquinoline sulfate, potassium hydroxyquinoline sulfate, hymexazol, cyazofamid, imazalil sulfate, imazalil, 5 imibenconazole, iminoctadine, iminoctadine triacetate, ipconazole, iprobenfos, iprodione, iprovalicarb, isoprothiolane, kasugamycin, kasugamycin hydrochloride hydrate, kresoxim-methyl, mancopper, mancozeb, maneb, mepanipyrim, mepronil, mercuric chloride, mercuric oxide, mercurous chloride, metalaxyl, metalaxyl-M, metam-sodium, metam, metconazole, methasulfocarb, methyl isothiocyanate, 10 metiram, metominostrobin, mildiomyacin, myclobutanil, nabam, nickel bis(dimethyldithiocarbamate), nitrothal-isopropyl, nuarimol, octhilinone, ofurace, oleic acid, oxadixyl, oxine-copper, oxpoconazole fumarate, oxycarboxin, pefurazoate, penconazole, pencycuron, pentachlorophenol, sodium pentachlorophenoxide, pentachlorophenyl laurate, phenylmercury acetate, sodium 2- 15 phenylphenoxide, 2-phenylphenol, phthalide, picoxystrobin, piperalin, polyoxinspolyoxin B, polyoxin, polyoxorim, probenazole, prochloraz, procymidone, propamocarb hydrochloride, propamocarb, propiconazole, propineb, prothioconazole, pyrazophos, pyributicarb, pyrifenoxy, pyrimethanil, pyroquilon, quinoxifen, quintozone, silthiofam, spiroxamine, sulfur, tar oils, tebuconazole, 20 tecnazene, tetraconazole, thiabendazole, thifluzamide, thiophanate-methyl, thiram, tolclofos-methyl, tolylfluanid, triadimefon, triadimenol, triazoxide, tricyclazole, tridemorph, trifloxystrobin, triflumizole, triforine, triticonazole, validamycin, vinclozolin, zineb, ziram, zoxamide, phosphorous acid, pyraclostrobin and simeconazole.

25 Preferably, fungicidal active ingredient (c) is selected from diethofencarb, hexaconazole, cyprodinil, tebuconazole and bromuconazole.

Where the third active ingredient (c) as defined above is present in the composition, this compound may be present in an amount of (a) : (b) : (c) weight 30 ratio of from 1 : 0.01 : 0.01 to 1 : 20 : 20; the ratios of compound (a) and compound (c) varying independently from each other. Preferably, the (a) : (b) : (c) weight ratio may be of from 1 : 0.05 : 0.05 to 1 : 10 : 10.

Following compositions may be cited to illustrate in a non-limited manner the 35 present invention : compound 1 with 2-butoxy-6-iodo-3-propyl-benzopyran-4-one, compound 1 with 2,6-dichloro-N-[[3-chloro-5-(trifluoromethyl)-2-

- pyridinyl)methyl}benzamide, compound 1 with (Z)-N-[α -(cyclopropylmethoxyimino)-2,3-difluoro-6-(trifluoromethyl)benzyl]-2-phenylacetamide, compound 1 with benthiavalicarb, compound 1 with chlorothalonil, compound 1 with copper hydroxide, compound 1 with copper oxychloride, compound 1 with copper sulfate, compound 1 with copper sulfate (tribasic), compound 1 with cuprous oxide, compound 1 with cymoxanil, compound 1 with diclomezine, compound 1 with dichlofluanid, compound 1 with dithianon, compound 1 with dimethomorph, compound 1 with dodine, compound 1 with ethaboxam, compound 1 with fenpiclonil, compound 1 with fentin, compound 1 with ferbam, compound 1 with fluazinam, compound 1 with fludioxonil, compound 1 with flusulfamide, compound 1 with guazatine, compound 1 with iminoctadine, compound 1 with mancopper, compound 1 with mancozeb, compound 1 with maneb, compound 1 with metiram, compound 1 with methasulfocarb, compound 1 with nabam, compound 1 with nickel bis(dimethyldithiocarbamate), compound 1 with iprovalicarb, compound 1 with oxine-copper, compound 1 with propamocarb, compound 1 with propineb, compound 1 with quinoxifen, compound 1 with sulfur, compound 1 with silthiofam, compound 1 with thiram, compound 1 with tolylfluanid, compound 1 with triazoxide, compound 1 with zineb, compound 1 with ziram, compound 1 with phosphorous acid, compound 1 with fosetyl-Al, compound 1 with chlozolate, compound 1 with iprodione, compound 1 with procymidone, compound 1 with vinclozolin, compound 1 with captafol, compound 1 with captan, compound 1 with folpet, compound 1 with thiochlorfenphim, compound 2 with 2-butoxy-6-iodo-3-propyl-benzopyran-4-one, compound 2 with 2,6-dichloro-N-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl)methyl}benzamide, compound 2 with (Z)-N-[α -(cyclopropylmethoxyimino)-2,3-difluoro-6-(trifluoromethyl)benzyl]-2-phenylacetamide, compound 2 with benthiavalicarb, compound 2 with chlorothalonil, compound 2 with copper hydroxide, compound 2 with copper oxychloride, compound 2 with copper sulfate, compound 2 with copper sulfate (tribasic), compound 2 with cuprous oxide, compound 2 with cymoxanil, compound 2 with diclomezine, compound 2 with dichlofluanid, compound 2 with dithianon, compound 2 with dimethomorph, compound 2 with dodine, compound 2 with ethaboxam, compound 2 with fenpiclonil, compound 2 with fentin, compound 2 with ferbam, compound 2 with fluazinam, compound 2 with fludioxonil, compound 2 with flusulfamide, compound 2 with guazatine, compound 2 with iminoctadine, compound 2 with mancopper, compound 2 with mancozeb, compound 2 with maneb, compound 2 with metiram, compound 2 with methasulfocarb, compound 2 with

nabam, compound 2 with nickel bis(dimethyldithiocarbamate), compound 2 with
 iprovalicarb, compound 2 with oxine-copper, compound 2 with propamocarb,
 compound 2 with propineb, compound 2 with quinoxifen, compound 2 with sulfur,
 compound 2 with silthiofam, compound 2 with thiram, compound 2 with
 5 tolylfluanid, compound 2 with triazoxide, compound 2 with zineb, compound 2 with
 ziram, compound 2 with phosphorous acid, compound 2 with fosetyl-Al, compound
 2 with chlozolate, compound 2 with iprodione, compound 2 with procymidone,
 compound 2 with vinclozolin, compound 2 with captafol, compound 2 with captan,
 compound 2 with folpet, compound 2 with thiochlorfenphim, compound 3 with 2-
 10 butoxy-6-iodo-3-propyl-benzopyran-4-one, compound 3 with 2,6-dichloro-N-[[3-
 chloro-5-(trifluoromethyl)-2-pyridinyl]methyl]benzamide, compound 3 with (Z)-N-
 [α -(cyclopropylmethoxyimino)-2,3-difluoro-6-(trifluoromethyl)benzyl]-2-
 phenylacetamide, compound 3 with benthiavalicarb, compound 3 with
 chlorothalonil, compound 3 with copper hydroxide, compound 3 with copper
 15 oxychloride, compound 3 with copper sulfate, compound 3 with copper sulfate
 (tribasic), compound 3 with cuprous oxide, compound 3 with cymoxanil, compound
 3 with diclomezine, compound 3 with dichlofluanid, compound 3 with dithianon,
 compound 3 with dimethomorph, compound 3 with dodine, compound 3 with
 ethaboxam, compound 3 with fenpiclonil, compound 3 with fentin, compound 3 with
 20 ferbam, compound 3 with fluazinam, compound 3 with fludioxonil, compound 3
 with flusulfamide, compound 3 with guazatine, compound 3 with iminoctadine,
 compound 3 with mancopper, compound 3 with mancozeb, compound 3 with maneb,
 compound 3 with metiram, compound 3 with methasulfocarb, compound 3 with
 nabam, compound 3 with nickel bis(dimethyldithiocarbamate), compound 3 with
 25 iprovalicarb, compound 3 with oxine-copper, compound 3 with propamocarb,
 compound 3 with propineb, compound 3 with quinoxifen, compound 3 with sulfur,
 compound 3 with silthiofam, compound 3 with thiram, compound 3 with
 tolylfluanid, compound 3 with triazoxide, compound 3 with zineb, compound 3 with
 ziram, compound 3 with phosphorous acid, compound 3 with fosetyl-Al, compound
 30 3 with chlozolate, compound 3 with iprodione, compound 3 with procymidone,
 compound 3 with vinclozolin, compound 3 with captafol, compound 3 with captan,
 compound 3 with folpet, compound 3 with thiochlorfenphim

The composition according to the present invention may further comprise an
 35 other additional component such as an agriculturally acceptable support, carrier or
 filler.

In the present specification, the term "support" denotes a natural or synthetic, organic or inorganic material with which the active material is combined to make it easier to apply, notably to the parts of the plant. This support is thus generally inert and should be agriculturally acceptable. The support may be a solid or a liquid.

- 5 Examples of suitable supports include clays, natural or synthetic silicates, silica, resins, waxes, solid fertilisers, water, alcohols, in particular butanol, organic solvents, mineral and plant oils and derivatives thereof. Mixtures of such supports may also be used.

10 The composition may also comprise other additional components. In particular, the composition may further comprise a surfactant. The surfactant can be an emulsifier, a dispersing agent or a wetting agent of ionic or non-ionic type or a mixture of such surfactants. Mention may be made, for example, of polyacrylic acid salts, lignosulphonic acid salts, phenolsulphonic or naphthalenesulphonic acid salts, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty
15 amines, substituted phenols (in particular alkylphenols or arylphenols), salts of sulphosuccinic acid esters, taurine derivatives (in particular alkyl taurates), phosphoric esters of polyoxyethylated alcohols or phenols, fatty acid esters of polyols, and derivatives of the above compounds containing sulphate, sulphonate and phosphate functions. The presence of at least one surfactant is generally essential
20 when the active material and/or the inert support are water-insoluble and when the vector agent for the application is water. Preferably, surfactant content may be comprised between 5% and 40% by weight of the composition.

Additional components may also be included, e.g. protective colloids, adhesives, thickeners, thixotropic agents, penetration agents, stabilisers, sequestering
25 agents. More generally, the active materials can be combined with any solid or liquid additive, which complies with the usual formulation techniques.

In general, the composition according to the invention may contain from 0.05 to 99% (by weight) of active material, preferably 10 to 70% by weight.

30 Compositions according to the present invention can be used in various forms such as aerosol dispenser, capsule suspension, cold fogging concentrate, dustable powder, emulsifiable concentrate, emulsion oil in water, emulsion water in oil, encapsulated granule, fine granule, flowable concentrate for seed treatment, gas (under pressure), gas generating product, granule, hot fogging concentrate, macrogranule, microgranule, oil dispersible powder, oil miscible flowable
35 concentrate, oil miscible liquid, paste, plant rodlet, powder for dry seed treatment, seed coated with a pesticide, soluble concentrate, soluble powder, solution for seed

treatment, suspension concentrate (flowable concentrate), ultra low volume (ulv) liquid, ultra low volume (ulv) suspension, water dispersible granules or tablets, water dispersible powder for slurry treatment, water soluble granules or tablets, water soluble powder for seed treatment and wettable powder.

5 These compositions include not only compositions which are ready to be applied to the plant or seed to be treated by means of a suitable device, such as a spraying or dusting device, but also concentrated commercial compositions which must be diluted before they are applied to the crop.

10 The fungicidal compositions of the present invention can be used to curatively or preventively control phytopathogenic fungi of crops. Thus, according to a further aspect of the present invention, there is provided a method for preventively or curatively controlling phytopathogenic fungi of crops characterised in that a fungicidal composition as hereinbefore defined is applied to the seed, the plant and/or to the fruit
15 of the plant or to the soil in which the plant is growing or in which it is desired to grow.

The composition as used against phytopathogenic fungi of crops comprises an effective and non-phytotoxic amount of an active material of general formula (I).

20 The expression "effective and non-phytotoxic amount" means an amount of composition according to the invention which is sufficient to control or destroy the fungi present or liable to appear on the crops, and which does not entail any appreciable symptom of phytotoxicity for the said crops. Such an amount can vary within a wide range depending on the fungus to be combated or controlled, the type of crop, the climatic conditions and the compounds included in the fungicidal composition according to the invention.

25 This amount can be determined by systematic field trials, which are within the capabilities of a person skilled in the art.

30 The method of treatment according to the present invention is useful to treat propagation material such as tubers or rhizomes, but also seeds, seedlings or seedlings pricking out and plants or plants pricking out. This method of treatment can also be useful to treat roots. The method of treatment according to the present invention can also be useful to treat the overground parts of the plant such as trunks, stems or stalks, leaves, flowers and fruits of the concerned plant.

35 Among the plants that can be protected by the method according to the invention, mention may be made of cotton; flax; vine; fruit crops such as *Rosaceae* sp. (for instance pip fruits such as apples and pears, but also stone fruits such as apricots, almonds and peaches), *Ribesioideae* sp., *Juglandaceae* sp., *Betulaceae* sp.,

Anacardiaceae sp., *Fagaceae* sp., *Moraceae* sp., *Oleaceae* sp., *Actinidaceae* sp., *Lauraceae* sp., *Musaceae* sp. (for instance banana trees and plantains), *Rubiaceae* sp., *Theaceae* sp., *Sterculiaceae* sp., *Rutaceae* sp. (for instance lemons, oranges and grapefruits); leguminous crops such as *Solanaceae* sp. (for instance tomatoes),
 5 *Liliaceae* sp., *Asteraceae* sp. (for instance lettuces), *Umbelliferae* sp., *Cruciferae* sp., *Chenopodiaceae* sp., *Cucurbitaceae* sp., *Papilionaceae* sp. (for instance peas), *Rosaceae* sp. (for instance strawberries); big crops such as *Graminae* sp. (for instance maize, cereals such as wheat, rice, barley and triticale), *Asteraceae* sp. (for instance sunflower), *Cruciferae* sp. (for instance colza), *Papilionaceae* sp. (for
 10 instance soja), *Solanaceae* sp. (for instance potatoes), *Chenopodiaceae* sp. (for instance beetroots); horticultural and forest crops; as well as genetically modified homologues of these crops.

Among the plants and the possible diseases of these plants protected by the method according to the present invention, mention may be made of :

- 15 - wheat, as regards controlling the following seed diseases: fusaria (*Microdochium nivale* and *Fusarium roseum*), stinking smut (*Tilletia caries*, *Tilletia controversa* or *Tilletia indica*), septoria disease (*Septoria nodorum*) and loose smut;
- wheat, as regards controlling the following diseases of the aerial parts of the plant: cereal eyespot (*Tapesia yallundae*, *Tapesia acuiformis*), take-all
 20 (*Gaeumannomyces graminis*), foot blight (*F. culmorum*, *F. graminearum*), black speck (*Rhizoctonia cerealis*), powdery mildew (*Erysiphe graminis forma specie tritici*), rusts (*Puccinia striiformis* and *Puccinia recondita*) and septoria diseases (*Septoria tritici* and *Septoria nodorum*);
- wheat and barley, as regards controlling bacterial and viral diseases, for
 25 example barley yellow mosaic;
- barley, as regards controlling the following seed diseases: net blotch (*Pyrenophora graminea*, *Pyrenophora teres* and *Cochliobolus sativus*), loose smut (*Ustilago nuda*) and fusaria (*Microdochium nivale* and *Fusarium roseum*);
- barley, as regards controlling the following diseases of the aerial parts of the
 30 plant: cereal eyespot (*Tapesia yallundae*), net blotch (*Pyrenophora teres* and *Cochliobolus sativus*), powdery mildew (*Erysiphe graminis forma specie hordei*), dwarf leaf rust (*Puccinia hordei*) and leaf blotch (*Rhynchosporium secalis*);
- potato, as regards controlling tuber diseases (in particular *Helminthosporium solani*, *Phoma tuberosa*, *Rhizoctonia solani*, *Fusarium solani*), mildew (*Phytophthora infestans*) and certain viruses (virus Y);
 35

- potato, as regards controlling the following foliage diseases: early blight (*Alternaria solani*), mildew (*Phytophthora infestans*);
- cotton, as regards controlling the following diseases of young plants grown from seeds: damping-off and collar rot (*Rhizoctonia solani*, *Fusarium oxysporum*) and
5 black root rot (*Thielaviopsis basicola*);
- protein yielding crops, for example peas, as regards controlling the following seed diseases: anthracnose (*Ascochyta pisi*, *Mycosphaerella pinodes*), fusaria (*Fusarium oxysporum*), grey mould (*Botrytis cinerea*) and mildew (*Peronospora pisi*);
- 10 - oil-bearing crops, for example rape, as regards controlling the following seed diseases: *Phoma lingam*, *Alternaria brassicae* and *Sclerotinia sclerotiorum*;
- corn, as regards controlling seed diseases: (*Rhizopus* sp., *Penicillium* sp., *Trichoderma* sp., *Aspergillus* sp., and *Gibberella fujikuroi*);
- flax, as regards controlling the seed disease: *Alternaria linicola*;
- 15 - forest trees, as regards controlling damping-off (*Fusarium oxysporum*, *Rhizoctonia solani*);
- rice, as regards controlling the following diseases of the aerial parts: blast disease (*Magnaporthe grisea*), bordered sheath spot (*Rhizoctonia solani*);
- leguminous crops, as regards controlling the following diseases of seeds or
20 of young plants grown from seeds: damping-off and collar rot (*Fusarium oxysporum*, *Fusarium roseum*, *Rhizoctonia solani*, *Pythium* sp.);
- leguminous crops, as regards controlling the following diseases of the aerial parts: grey mould (*Botrytis* sp.), powdery mildews (in particular *Erysiphe cichoracearum*, *Sphaerotheca fuliginea* and *Leveillula taurica*), fusaria (*Fusarium oxysporum*, *Fusarium roseum*), leaf spot (*Cladosporium* sp.), alternaria leaf spot
25 (*Alternaria* sp.), anthracnose (*Colletotrichum* sp.), septoria leaf spot (*Septoria* sp.), black speck (*Rhizoctonia solani*), mildews (for example *Bremia lactucae*, *Peronospora* sp., *Pseudoperonospora* sp., *Phytophthora* sp.);
- fruit trees, as regards diseases of the aerial parts: monilia disease (*Monilia fructigenae*, *M. laxa*), scab (*Venturia inaequalis*), powdery mildew (*Podosphaera leucotricha*);
- 30 - vine, as regards diseases of the foliage: in particular grey mould (*Botrytis cinerea*), powdery mildew (*Uncinula necator*), black rot (*Guignardia biwelli*) and mildew (*Plasmopara viticola*);

- beetroot, as regards the following diseases of the aerial parts: cercospora blight (*Cercospora beticola*), powdery mildew (*Erysiphe beticola*), leaf spot (*Ramularia beticola*).

5 The fungicidal composition according to the present invention may also be used against fungal diseases liable to grow on or inside timber. The term "timber" means all types of species of wood, and all types of working of this wood intended for construction, for example solid wood, high-density wood, laminated wood, and plywood. The method for treating timber according to the invention mainly consists
10 in contacting one or more compounds of the present invention, or a composition according to the invention; this includes for example direct application, spraying, dipping, injection or any other suitable means.

15 The fungicidal composition according to the present invention may also be used in the treatment of genetically modified organisms with the compounds according to the invention or the agrochemical compositions according to the invention. Genetically modified plants are plants into in which genome a heterologous gene encoding a protein of interest has been stably integrated. The expression "heterologous gene encoding a protein of interest" essentially means
20 genes which give the transformed plant new agronomic properties, or genes for improving the agronomic quality of the transformed plant.


25 The dose of active material usually applied in the treatment according to the present invention is generally and advantageously between 10 and 2000 g/ha, preferably between 20 and 1500 g/ha for applications in foliar treatment. The dose of active substance applied is generally and advantageously between 1 and 200 g per 100 kg of seed, preferably between 2 and 150 g per 100 kg of seed in the case of seed treatment. It is clearly understood that the doses indicated above are given as illustrative examples of the invention. A person skilled in the art will know how to
30 adapt the application doses according to the nature of the crop to be treated.

35 The compositions according to the present invention may also be used fore the preparation of composition useful to curatively or preventively treat human and animal fungal diseases such as, for example, mycoses, dermatoses, trichophyton diseases and candidiases or diseases caused by *Aspergillus* spp. or *Candida* spp., for example *Aspergillus fumigatus* or *Candida albicans* respectively.

CLAIMS

5 1. A composition comprising :

a) a pyridylethylbenzamide derivative of general formula (I)



in which :

- p is an integer equal to 1, 2, 3 or 4;
- q is an integer equal to 1, 2, 3, 4 or 5;
- 10 - each substituent X is chosen, independently of the others, as being halogen, alkyl or haloalkyl;
- each substituent Y is chosen, independently of the others, as being halogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, amino, phenoxy, alkylthio, dialkylamino, acyl, cyano, ester, hydroxy, aminoalkyl, benzyl, haloalkoxy, halosulphonyl, halothioalkyl,
- 15 alkoxyalkenyl, alkylsulphonamide, nitro, alkylsulphonyl, phenylsulphonyl or benzylsulphonyl;
- as to the N-oxides of 2-pyridine thereof;
- and
- b) a compound capable of inhibiting the spores germination or mycelium growth by
- 20 acting on different metabolic routes;
- in a (a) / (b) weight ratio of from 0.01 to 20.

2. A composition according to claim 1, characterised in that p is 2.

25 3. A composition according to claim 1 or 2, characterised in that q is or 2.

4. A composition according to any of the claims 1 to 3, characterised in that X is chosen, independently of the others, as being halogen or haloalkyl.

5. A composition according to any of the claims 1 to 4, characterised in that X is chosen independently of the others, as being a chlorine atom or a trifluoromethyl group.
- 5 6. A composition according to any of the claims 1 to 5, characterised in that Y is chosen, independently of the others, as being halogen or haloalkyl.
7. A composition according to any of the claims 1 to 6, characterised in that Y is chosen, independently of the others, as being a chlorine atom or a trifluoromethyl group.
- 10 8. A composition according to any of the claims 1 to 7, characterised in that the compound of general formula (I) is :
- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide;
15 - N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-iodobenzamide; or
- N-{2-[3,5-dichloro-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide .
9. A composition according to claim 8, characterised in that the compound of general formula (I) is N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide.
- 20 10. A composition according to any of the claims 1 to 9, characterised in that the compound capable of inhibiting the spores germination or mycelium growth by acting on different metabolic routes is a dicarboximide derivative.
- 25 11. A composition according to claim 10, characterised in that the dicarboximide derivative is chlozolate, iprodione, procymidone or vinclozolin.
12. A composition according to any of the claims 1 to 9, characterised in that the compound capable of inhibiting the spores germination or mycelium growth by acting on different metabolic routes is a phthalimide derivative.
- 30 13. A composition according to claim 12, characterised in that the phthalimide derivative is captan, captan, folpet or thiochlorfenphim.
- 35

14. A composition according to any of the claims 1 to 9, characterised in that the compound capable of inhibiting the spores germination or mycelium growth by acting on different metabolic routes is 2-butoxy-6-iodo-3-propyl-benzopyran-4-one, 2,6-dichloro-N-{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl}benzamide, (Z)-
 5 N-[α -(cyclopropylmethoxyimino)-2,3-difluoro-6-(trifluoromethyl)benzyl]-2-phenylacetamide, benthiavalicarb, chlorothalonil, copper hydroxide, copper oxychloride, copper sulfate, copper sulfate (tribasic), cuprous oxide, cymoxanil, diclomezine, dichlofluanid, dithianon, dimethomorph, dodine, ethaboxam, fencpiclonil, fentin, ferbam, fluazinam, fludioxonil, flusulfamide, guazatine,
 10 iminoctadine, mancozeb, mancozeb, maneb, metiram, methasulfocarb, nabam, nickel bis(dimethyldithiocarbamate), iprovalicarb, oxine-copper, propamocarb, propineb, quinoxifen, sulfur, silthiofam, thiram, tolylfluanid, triazoxide, zineb, ziram, phosphorous acid or fosetyl-Al.

15

15. A composition according to any one of the claims 1 to 14 further comprising a fungicidal compound (c).

20

16. A composition according to claim 15, characterised in that the fungicidal compound (c) is selected from diethofencarb, hexaconazole, cyprodinil, tebuconazole and bromuconazole.

25

17. A composition according to any one of the claims 1 to 16, characterised in that it further comprises an agriculturally acceptable support, carrier, filler and/or surfactant.

30

18. A method for preventively or curatively controlling phytopathogenic fungi of crops, characterised in that an effective and non-phytotoxic amount of a composition according to any one of the claims 1 to 17 is applied to the seed, the plant and/or to the fruit of the plant or to the soil in which the plant is growing or in which it is desired to grow.

ABSTRACT

5 A composition comprising at least a pyridylethylbenzamide derivative of general formula (I) (a) and a compound capable of inhibiting the spores germination or mycelium growth by acting on different metabolic routes (b) in a (a) / (b) weight ratio of from 0.01 to 20.

10 A composition further comprising an additional fungicidal compound.

A method for preventively or curatively combating the phytopathogenic fungi of crops by using this composition.

